



Irritant sprays: clinical effects and management

Recommendations for Healthcare Professionals (Forensic Physicians, Custody Nurses and Paramedics)

Dec 2017 Review date Dec 2020 – check www.fflm.ac.uk for latest update

The medico-legal guidelines and recommendations published by the Faculty are for general information only. Appropriate specific advice should be sought from your medical defence organisation or professional association. The Faculty has one or more senior representatives of the MDOs on its Board, but for the avoidance of doubt, endorsement of the medico-legal guidelines or recommendations published by the Faculty has not been sought from any of the medical defence organisations.

Introduction

CS & PAVA irritant spray effects and management

Irritant (formerly known as ‘incapacitant sprays’) augment the range of ‘less-lethal’ tactical options available to police officers confronted by potentially aggressive or violent individuals or those with acute behavioural disturbance.

The terminology for CS and PAVA sprays used by police personnel was changed in the UK as the sprays are not ‘incapacitants’ in the way that word is defined in the Chemical Weapon Convention; they are ‘Riot Control Agents’ (i.e. their effects can be reversed without medical intervention). The Home Office Centre for Applied Science and Technology recommended the adoption of the term ‘irritant’ rather than ‘incapacitant’ for police CS and PAVA Sprays. This terminology will be used in these recommendations.

The agents currently used by police services in the UK are:

- CS
- PAVA

Both are contained in a suitable solvent as described in the Home Office Standard for Police Chemical Irritant Sprays.

They are intended to be used to spray the face of a person at a distance of up to 3-4 metres, delivering the active chemical to the eyes, nose and mouth. This causes irritation to the eyes, upper respiratory tract and skin.

In most cases the symptoms and signs are short-lived, resolving over 15 to 30 minutes and requiring little or no medical intervention although some individuals may experience effects for 2.5 hours or more. Longer exposure or exposure in confined spaces may result in enhanced or prolonged effects.

The broad principles of effect, treatment and management are the same for both CS spray and PAVA spray. No reported clinical studies have compared the nature and incidence of their different effects.

Chemical irritants can cause severe injury, permanent disabilities, and in rare cases, death. The true incidence of morbidity (and possible mortality) of irritant spray remains unknown in the absence of prospective clinical studies of appropriate statistical power.

CS Irritant spray

Chemical name	CS O-chlorbenzylidene malonitrile, 5% solution
Solvent	Methyl Isobutyl Ketone (MIBK)
Propellant	Nitrogen for hand held spray
Formulation used	Liquid Spray

CS is a solid at room temperature but is dissolved in an organic solvent to be used as a liquid aerosol. The solvent evaporates leaving the CS particles to give their effects.

PAVA Irritant spray

Chemical name	Pelargonic acid vanillylamide, 0.3% solution (nonivamide)
Solvent	Monopropylene glycol, ethanol, water
Propellant	Nitrogen
Formulation used	Liquid spray

PAVA is a synthetic variant of capsaicin (the active ingredient of natural pepper).

CS and PAVA fail to affect about 10% of people exposed to spray. The reasons for this are varied and may include intoxication with other drugs or alcohol; mental health issues; or other causes of acute behavioural disturbance. Agitation may impede initial decontamination and increase the risks of secondary contamination of self or others.



General advice

In the majority of cases the main effects pass off after 15 to 30 minutes and medical intervention is not required, although skin and eye signs and symptoms may persist beyond 2.5 hours.

It is very important to reassure the affected individual.

Restraint in the prone position must be avoided after exposure. If restrained, breathing must be monitored constantly.

Any effects that last for > 6 hours should generally be referred for relevant specialist assessment.

- The most important action is to stop continued exposure by removal of the affected individual from the contaminated environment to a well-ventilated area, preferably with a free flow of air (enhanced by the use of electric fans), and careful removal of contaminated clothing (which should be placed in sealed plastic bags).
- Water should not be used at an early stage to attempt to remove residue as it will exacerbate symptoms or cause symptom relapse.
- Avoid rubbing the eyes. If an individual cannot open their eyes after 15-20 minutes, then copious amounts of cool tap water should be used to flush remaining irritant from the skin of the face (sterile normal saline should be used to treat eyes – see below). Contact lenses should be removed as soon as possible. Depending on type they may not be reusable. Under no circumstances should warm water be used to irrigate as this can reactivate the irritants.
- Those dealing with the contaminated individual should wear gloves and eye protection to avoid cross contamination.
- Care should be taken to avoid CS entering air conditioning or ventilation systems.
- Each individual should be fully examined by an appropriately skilled and trained healthcare professional, with particular reference to eyes, oral and nasal cavity, respiratory system and skin.
- Subjects with any signs or symptoms that do not resolve after 6 hours should be referred for hospital assessment.
- Persons sprayed or contaminated with CS or PAVA should be advised to contact their general practitioner or attend their emergency department with a copy of these guidelines if problems develop once they have been released from custody, or complete resolution of signs and symptoms does not occur.
- Normal machine washing will decontaminate clothing but it may take several washes to be fully successful.

Specific effects and management

The need for specialist referral will be determined by a full medical assessment, based on the existing signs and symptoms and not on the irritant agent used.

Eyes

Clinical effects – generally expected duration of most intense effects

- lachrymation (tears) (<15 mins);
- pain (<30 mins);
- blepharospasm (eyelids closed) (<30 mins);
- conjunctival erythema (redness) (<30 mins);
- reduced visual acuity (blurred vision) (<30 mins);
- photophobia (sensitivity to light) (<60 mins);
- periorbital oedema (swelling around the eye);
- damage to the ocular surface from the direct trauma of a high-pressure jet;
- iritis may develop as a non-specific response
- conjunctivitis;
- corneal abrasions due to rubbing the eyes.

Management

- air can be blown with a fan directly onto the eyes to encourage evaporation;
- or exposure to external air/wind;
- if eye symptoms persist for more than one hour irrigate eyes with sterile normal saline solution (this may temporarily exacerbate symptoms as the vapour passes into solution) prior to hospital referral;
- contact lenses should be removed and either discarded (soft) or cleaned with 10 washes and soaks. It may take several weeks for the eye to settle down enough to allow a return to contact lens wear (persistent symptoms must be reviewed by an optometrist);
- prophylactic antibiotics have no role;
- if eye symptoms do not resolve after 6 hours or corneal abrasion is identified refer for formal ophthalmic assessment.

Mouth

Clinical effects

- stinging or burning sensation;
- possible nausea and vomiting (rare).

Management

- symptomatic, based on clinical findings.



Respiratory tract

Clinical effects

- nose discomfort, pain & rhinorrhoea (<30 mins);
- sneezing & coughing;
- sore throat;
- shortness of breath;
- bronchospasm (rare);
- laryngospasm (rare);
- tracheitis;
- bronchitis (rare);
- pulmonary oedema may develop 12 to 24 hours after excessive exposure (rare);
- NB Patients with pre-existing respiratory disease, such as asthma or bronchitis, are more at risk of severe effects.

Management

- The majority of respiratory tract symptoms and signs (e.g. dyspnoea, chest tightness and irregular breathing) should settle within 15 minutes of the exposure;
- If there is evidence of bronchospasm that does not respond to bronchodilator refer to hospital for assessment and treatment. Humidified oxygen may provide some relief;
- Persistent respiratory symptoms necessitate review by a respiratory physician.

Skin

Clinical effects

- burning sensation & erythema (<24 hrs);
- chemical burns, blistering;
- allergic contact dermatitis (rare - but if in law enforcement personnel regularly exposed to irritant spray may require changes in work practice – referral to Occupational Health teams should be made);
- leukoderma (rare);
- initiation or exacerbation of seborrhoeic dermatitis (rare);
- aggravation of rosacea (rare).

Management

- exposure to air and fan;
- exposure to fresh air will normally result in a significant recovery within 15 – 20 minutes;
- if reactions do persist beyond this period then copious amounts of cool tap water should be used to flush remaining irritant from the face and skin;
- under no circumstances should warm water be used as this can reactivate irritants;
- treat chemical burns as thermal burns;
- topical steroids can be used for contact dermatitis;

- delayed skin irritation (due to MIBK), occurring 8 to 16 hours after exposure, is seen in a significant number; these symptoms can take up to one week to resolve.
- persistence of new skin conditions or aggravation of chronic conditions beyond 48 hours should be assessed by the person's GP and referral to a dermatologist may be required.

Cardiovascular effects

Clinical effects

- Pre-existing cardiac problems can be worsened and hypertension exacerbated after exposure. For example, angina attacks may develop.

Management

- Symptomatic treatment e.g. glyceryl trinitrate;
- Refer to hospital if any concerns at examination (e.g. persistent tachycardia, arrhythmias, hypertension, hypotension).

Other

Psychological effects

- In one study, one quarter of those exposed to CS spray were diagnosed with Post Traumatic Stress Disorder; a past psychiatric history and a more external locus of control was associated with post-traumatic morbidity.

Management

- Consideration should be given to early psychological intervention if the individual is perceived to be at risk.



References

- Association of Chief Police Officer of England, Wales & Northern Ireland
Guidance on the Use of Incapacitant Spray
 ACPO 2009
- Carron P-N, Yersin B
Management of the effects of exposure to tear gas
 BMJ 2009; 338: 1554-8
- Croft S
HOSDB Standard for CS and PAVA Sprays for Operational Police Use
 2008. Revision 1. 38/08
- Euripidou E, MacLehose R, Fletcher A
An investigation into the short term and medium term health impacts of personal incapacitant sprays. A follow up of patients reported to the National Poisons Information Service
 Emerg. Med. J 2004;21;548-552
- Haar RJ, Iacopino V, Ranadive N, Weiser SD, Dandu M
Health impacts of chemical irritants used for crowd control: a systematic review of the injuries and deaths caused by tear gas and pepper spray
 BMC Public Health 2017 Oct 19;17(1):831
 doi: 10.1186/s12889-017-4814-6
- Karagama Y, Newton J, Newbegin C
Short-term and long-term physical effects of exposure to CS spray
 R Soc Med 2003; 96: 172-4
- Nathan R, Wood H, Rix K, Wright E
Long-term psychiatric morbidity in the aftermath of CS spray trauma
 Med Sci Law 2003 Apr;43(2):98-104
- Payne-James JJ, Smith G, Rivers E, O'Rourke S, Stark M, Sutcliffe N
Effects of incapacitant spray deployed in the restraint and arrest of detainees in the Metropolitan Police Service area, London, UK: a prospective study
 Forensic Sci Med Pathol. 2014 Mar;10(1):62-8. doi: 10.1007/s12024-013-9494-7. Epub 2013 Nov 10
- Payne-James JJ, Rivers E, Green P, Johnston A
Trends in less-lethal use of force techniques by police services within England and Wales: 2007-2011
 Forensic Sci Med Pathol. 2014 Mar;10(1):50-5
 doi: 10.1007/s12024-013-9492-9. Epub 2013 Oct 31
- Payne-James JJ. Irritant Spray. In: Payne-James JJ, Byard RW (eds)
Encyclopedia of Forensic & Legal Medicine, 2nd Edition
 Elsevier 2015
- Rotherham Doncaster and South Humber NHS Foundation Trust
Guidelines For Staff In The Event That Police Use A Taser Or Irritant Spray On An Inpatient Whilst Responding To An Incident
 2015 p. 8-10
- Southward R D
CS incapacitant spray
 J Accid Emerg Med 2000; 17:76
- Weir E
The health impact of crowd control agents
 Canadian Medical Association Journal 2001; 26: 164(13): 1889-1890
- Worthington E, Nee Patrick A
CS exposure – clinical effects and management
 J Accid Emerg Med 1999; 16:168-170